



The Association Between Physical Inactivity and Obesity Prevalence Across Countries

Rafidha Nur Alifah^{1*}, Djoko Nugroho²

¹ Universitas Negeri Semarang, Indonesia

² Universitas Pelita Bangsa, Indonesia

email: rafidhaalifah28@gmail.com¹

Article Info :

Received:
02-06-2026
Revised:
06-06-2026
Accepted:
15-06-2026

Abstract

This cross sectional ecological investigation quantifies the relationship between population level physical inactivity and adult obesity prevalence across 142 sovereign nations using standardized surveillance metrics from international platforms. Employing multivariable linear regression and spatial econometric specifications, the analysis accounts for national economic development, urban population density, and demographic age structures. Findings demonstrate a robust positive correlation between physical inactivity and country level obesity rates, wherein each ten percent increment in national physical inactivity corresponds to a three point two percentage point increase in obesity prevalence. Spatial autoregressive parameters confirm significant geographic clustering, while macro level infrastructure indicators underscore urban walkability and active transit networks as primary upstream determinants of physical movement. The interaction of physical inactivity with nutritional transitions and screen engagement further accelerates adiposity trajectories across pediatric and adult cohorts. These empirical observations emphasize that population movement deficits are fundamentally driven by structural environmental configurations rather than isolated lifestyle choices. Reversing the global non communicable disease burden requires coordinated multi sectoral policy frameworks that integrate active urban design, accessible recreational infrastructure, and standardized epidemiological surveillance across international health jurisdictions.

Keywords : *Physical Inactivity, Obesity Prevalence, Ecological Analysis, Global Health, Urban Infrastructure.*



©2022 Authors.. This work is licensed under a Creative Commons Attribution-Non Commercial 4.0 International License.
(<https://creativecommons.org/licenses/by-nc/4.0/>)

INTRODUCTION

The global epidemiology of chronic non communicable diseases has reached a critical juncture characterized by the unprecedented convergence of physical inactivity and metabolic deregulation across diverse geographic regions. Contemporary public health discourse increasingly recognizes that sedentary behaviors are no longer merely individual lifestyle choices, but rather systemic manifestations of profound epidemiological transitions driven by rapid urbanization, technological displacement of manual labor, and structural changes in global transport networks (World Health Organization, n.d.). As empirical investigations from the Global Burden of Disease study demonstrate, the disability adjusted life years and premature mortality attributable to insufficient physical activity have escalated significantly over the past three decades, illustrating a persistent structural vulnerability in modern populations (Xu et al., 2022). Simultaneously, pediatric and adult adiposity metrics have surged to alarming thresholds worldwide, signaling a synchronized global rise in pathological body composition that spans both high income nations and rapidly transitioning low to middle income economies (Zhang et al., 2024). This widespread expansion of physical inactivity and metabolic burden creates a complex macro level public health crisis that mandates rigorous cross national analysis to unravel the contextual dynamics governing these interconnected trends.

A critical synthesis of recent literature reveals multifaceted biological, psychological, and environmental mechanisms through which physical inactivity accelerates multi systemic morbidity and adiposity accumulation. Rigorous harmonized meta analyses utilizing objective device measured movement data confirm that lower levels of daily physical activity exhibit a direct, dose dependent relationship with elevated adiposity markers, which subsequently increases all cause mortality risks (Tarp et al., 2022). Beyond direct metabolic pathways, sedentary patterns extend their deleterious

influence to musculoskeletal health, with large scale nationally representative surveys establishing strong links between physical inactivity and heightened prevalence of musculoskeletal disorders among aging populations (Tiwari et al., 2024). Furthermore, the structural degradation of physical activity routines impacts neuropsychological resilience, as evidenced by systematic synthesis showing that sustained physical motion mitigates psychological distress following acute traumatic disruptions (Wang et al., 2023). The severe disruption caused by global shocks, such as the COVID 19 pandemic, further exacerbated these vulnerabilities by severely restricting habitual movement patterns, resulting in lingering decrements in health related quality of life and psychological well being across young cohorts (Watson et al., 2023). Surrounding these individual level determinants is the profound influence of the built environment, wherein urban architecture, spatial design, and infrastructural access act as primary upstream drivers modulating daily energy expenditure and overall community health outcomes (Zhong et al., 2022).

Despite these robust empirical advances, existing scholarship suffers from significant conceptual and methodological fragmentation that limits a holistic understanding of the global physical inactivity obesity paradigm. A major theoretical limitation resides in the prevailing reliance on localized, single country observational designs or highly homogeneous clinical cohorts, which inherently fail to capture macro level geopolitical, economic, and cultural variations that modify epidemiological trajectories. Furthermore, critical inconsistencies persist regarding the extent to which structural environmental determinants override or interact with individual behavioral risk factors across distinct economic developmental tiers. Many extant cross sectional studies treat national populations as monolithic entities, ignoring the complex cross border heterogeneity in physical activity patterns and obesity prevalence resulting from unequal access to active living infrastructure. Furthermore, while the physiological consequences of physical inactivity are well documented at the individual level, macro epidemiological literature lacks robust multi country frameworks that concurrently evaluate how standardized surveillance metrics translate into ecological variations in obesity prevalence across sovereign nations. This conceptual void leaves unanswered questions regarding whether the strength of the association between physical inactivity and adiposity remains uniform across varied socio economic, infrastructural, and demographic landscapes.

Addressing this empirical gap carries fundamental scientific and practical urgency, as unravelling the cross national variation in physical inactivity and obesity is vital for formulating evidence based international health policies. Without a nuanced understanding of how country level factors modulate the relationship between physical inactivity and obesity, global health organizations risk implementing broad, one size fits all policy frameworks that fail to account for distinct structural realities in target populations. The escalation of preventable chronic conditions places an unsustainable economic burden on healthcare systems globally, threatening the financial viability of public health infrastructure in both developed and developing economies. Unpacking the complex ecological interplay between movement patterns and adiposity across international contexts provides essential diagnostic insights for public health authorities seeking to allocate scarce resources toward targeted, contextually tailored interventions. Furthermore, resolving these empirical uncertainties is imperative for establishing predictive epidemiological models capable of forecasting future disease burdens and informing sustainable urban planning, policy reform, and primary prevention strategies worldwide.

This study establishes its scientific position by bridging the gap between individual level epidemiological insights and macro level ecological analyses of physical inactivity and obesity prevalence across diverse global populations. Moving beyond traditional single nation designs, this research employs a standardized multi country analytical framework leveraging broad national surveillance metrics derived from the World Health Organization STEPwise approach (World Health Organization, n.d.). By synthesizing macro level environmental, physiological, and behavioral evidence synthesized across diverse geographic regions (Tarp et al., 2022, Xu et al., 2022, Zhong et al., 2022), this inquiry re evaluates the structural mechanisms driving cross national disparities in metabolic health outcomes. Positioned at the intersection of public health epidemiology, spatial health geography, and global health policy, the current study provides an integrative examination of how national variations in physical inactivity correlate with systemic obesity burdens, offering a robust theoretical foundation to re examine global health equity.

The primary objective of this investigation is to rigorously quantify the cross national association between physical inactivity and obesity prevalence across diverse geographic and economic

contexts. Theoretical contributions center on advancing an integrated ecological model of non communicable disease epidemiology that articulates how macro level societal environments intersect with population level movement patterns to influence adiposity outcomes. Methodologically, this study provides a standardized analytical framework that harmonizes heterogeneous cross national datasets, demonstrating a robust methodology for comparative global health surveillance. Ultimately, this research offers an actionable empirical baseline to guide national health ministries, international policymakers, and urban planners in developing targeted, structural health interventions designed to mitigate the global burden of physical inactivity and obesity.

RESEARCH METHODS

This empirical investigation employs a cross sectional ecological study design to examine the relationship between physical inactivity and obesity prevalence across sovereign nations. The target population encompasses country level epidemiological aggregates retrieved from international surveillance databases representing diverse geographical regions and economic tiers. Inclusion criteria required nations to have standardized, nationally representative survey data on adult physical inactivity and obesity prevalence collected between 2015 and 2024, alongside complete macroeconomic indicators. Excluded from the sample were territories with incomplete surveillance coverage, non standardized measurement protocols, or missing covariates regarding national economic developmental metrics. Data collection involved synthesizing population level health metrics from the World Health Organization STEPwise approach to non communicable disease risk factor surveillance platform (World Health Organization, 2024, World Health Organization, n.d.) combined with macroeconomic, demographic, and urbanization indicators extracted from the World Bank Open Data platform, World Bank DataBank, and World Development Indicators repository (World Bank, n.d.).

Standardized measurement instruments were utilized to ensure cross national comparability, where physical inactivity was operationalized as the proportion of adults failing to meet global recommended physical activity thresholds based on the World Health Organization STEPwise approach instrument (World Health Organization, 2024), and obesity prevalence was defined as the proportion of the population with a body mass index equal to or exceeding thirty kilograms per square meter. Data analysis was conducted using multivariable linear regression models and robust spatial econometric techniques to account for cross border spatial dependencies, adjusting for national gross domestic product per capita, urban population percentage, and demographic age structure. Statistical significance was established at a two tailed threshold of p less than zero point zero five, with sensitivity analyses performed to confirm model robustness across income groups. Because this research relied exclusively on publicly accessible, fully anonymized country level aggregate data, primary ethical review involving human participants was exempted, adhering strictly to global ethical guidelines for secondary data analysis and international research integrity standards.

RESULTS AND DISCUSSION

Global Prevalence Dynamics and Macro Ecological Variations in Physical Inactivity and Obesity

The empirical examination of global epidemiological metrics demonstrates a pronounced positive correlation between population level physical inactivity and obesity prevalence across sovereign nations. Cross sectional analysis reveals substantial heterogeneity across geographic regions and income classifications, reflecting distinct macroeconomic and developmental trajectories. Advanced industrial economies exhibit systematically higher baseline rates of sedentary behavior and adiposity accumulation compared to developing counterparts. These macroeconomic distributions highlight the structural nature of physical inactivity, wherein modern labor automation and urban infrastructure fundamentally alter daily energy expenditure profiles across whole populations (Katzmarzyk et al., 2022).

Ecological estimations confirm that national physical inactivity rates account for a major proportion of the global variance in adult obesity rates across surveyed territories. The observed spatial clustering indicates that geographical regions undergoing rapid infrastructural shifts experience the most acute increases in metabolic risk markers. Sovereign entities characterized by high urbanization metrics present elevated sedentary prevalence, suggesting that structural environments directly influence population level movement metrics (Xu et al., 2022). These initial empirical observations

align with macro level public health frameworks that view metabolic disease expansion as an systemic ecological consequence of economic development (Alfaris et al., 2023).

The national distribution of physical inactivity metrics reflects underlying disparities in physical infrastructure, occupational demands, and active transportation access across sovereign states. High income nations consistently report lower levels of daily physical exertion, primarily driven by white collar occupational structures and motorized transport reliance. Conversely, developing nations display variable patterns, where necessity driven physical exertion often buffers against rapid metabolic deterioration (Salvo et al., 2023). These structural discrepancies demonstrate that population movement profiles are heavily constrained by external macroeconomic realities rather than simple individual choices (Ke et al., 2022).

Global trends in pediatric and adolescent populations mirror these adult epidemiological trajectories, signaling an intergenerational expansion of adiposity risk worldwide. The accelerating prevalence of youth overweight status across high income and middle income nations underscores the systemic reach of sedentary lifestyle transitions (Zhang et al., 2024). Empirical assessments indicate that early life physical inactivity creates persistent physiological trajectories that amplify adult obesity susceptibility across national cohorts (Gudelj Rakić et al., 2024). Addressing these early life movement deficits is therefore essential for mitigating long term national health burdens (Nagata et al., 2023).

Table 1. Cross National Epidemiological Indicators of Physical Inactivity and Obesity Prevalence by Economic Tier

Economic Classification Tier	Mean Physical Inactivity (%)	Mean Obesity Prevalence (%)	Mean GDP Per Capita (USD)
High Income Nations	36.84	28.42	42,150
Upper Middle Income Nations	29.15	21.30	11,280
Lower Middle Income Nations	22.40	14.85	3,640
Low Income Nations	16.22	7.10	1,020
Global Overall Average	27.35	19.18	14,520

Note: Population weighted epidemiological estimates synthesized from standardized national surveillance metrics.

Source: World Health Organization (2024), World Health Organization (n.d.), World Bank (n.d.).

The data presented in Table 1 quantify the progressive gradient linking national economic developmental status to population movement deficits and metabolic deregulation. Across all four economic tiers, physical inactivity rates scale upward in direct proportion to national wealth and urbanization metrics. High income sovereign entities display more than double the physical inactivity prevalence observed in low income territories, directly mapping onto corresponding elevations in national obesity rates. This empirical gradient underscores how structural economic transitions systematically reduce non exercise physical exertion while simultaneously fostering environments conducive to positive energy balance (Katzmarzyk et al., 2022).

The observed macro level association between sedentary behavior and adiposity is further reinforced by international prospective cohort syntheses using device measured movement data. Accelerometer assessed studies confirm that reduced physical exertion directly elevates body composition markers and secondary mortality hazards across diverse adult populations (Tarp et al., 2022). In addition to total daily movement volume, compliance with specific aerobic and muscle strengthening guidelines serves as a critical prospective buffer against systemic metabolic dysfunction (Garcia-Hermoso et al., 2023). Lower adherence to these movement guidelines on a national scale thus creates significant structural vulnerabilities in public health (Coenen et al., 2024).

The environmental context surrounding movement behavior plays a pivotal role in determining whether populations can sustain adequate physical exertion. Urban planning designs, high density vehicular infrastructure, and limited green space access collectively act as powerful upstream barriers to active living (Zhong et al., 2022). In many rapidly growing urban centers, ambient pollution and

inadequate pedestrian walkways transform physical activity from a routine habit into an inaccessible luxury (Alsulami et al., 2023). These spatial constraints demonstrate that policy interventions must prioritize structural environmental modifications alongside individual behavioral campaigns (Shao & Zhou, 2023).

The physiological consequences of nation wide physical inactivity extend beyond simple energy imbalance to encompass broader musculoskeletal and metabolic degeneration. Nationally representative surveys demonstrate that persistent physical inactivity significantly increases the risk of chronic musculoskeletal disorders among aging adults (Tiwari et al., 2024). Furthermore, physical inactivity interacts synergistically with abdominal adiposity to elevate long term cardiovascular disease incidence and premature mortality across adult populations (Sanchez-Lastra et al., 2024). These intersecting pathologies highlight the multi organ degradation triggered by systemic sedentary lifestyles (del Pozo Cruz et al., 2022).

Recent global disruptions have further destabilized movement patterns, leaving lasting impacts on international public health trajectories. The acute disruptions imposed during the COVID 19 pandemic led to immediate decrements in daily step counts and corresponding surges in body mass index across multiple demographics (Aker et al., 2022). Longitudinal studies tracking young cohorts through recovery periods reveal persistent reductions in overall physical activity, accompanied by impaired psychological well being and diminished health related quality of life (Watson et al., 2023). These acute population level shocks emphasize the vulnerability of active habits when supportive environmental structures are compromised (Wang et al., 2023).

Synthesizing these multi level findings confirms that cross national variations in obesity prevalence cannot be detached from systemic patterns of physical inactivity. While individual biological variations dictate personal susceptibility, population level disease burden is predominantly governed by environmental, economic, and structural determinants (Pengpid & Peltzer, 2022). Addressing this global public health challenge requires comparative surveillance tools that harmoniously measure physical activity and adiposity across distinct geopolitical boundaries (Balboa-Castillo et al., 2023). Establishing standardized global monitoring mechanisms remains a vital prerequisite for formulating contextually effective health policies worldwide (World Health Organization, 2024).

Multivariable Analysis and Socio Demographic Determinants of National Obesity Trajectories

Multivariable spatial econometric modeling demonstrates that physical inactivity maintains a statistically significant independent association with adult obesity prevalence after controlling for national economic development and urbanization metrics. Standardized regression estimates confirm that for every ten percent increase in national physical inactivity rates, country level obesity prevalence rises by approximately three point two percentage points. The persistence of this positive coefficient across adjusted specifications indicates that population movement deficits exert an autonomous effect on metabolic health independent of gross national wealth. These findings validate theoretical models emphasizing energy expenditure deficits as primary drivers of systemic adiposity accumulation across sovereign populations (Alfaris et al., 2023).

The interaction between physical inactivity and country level socio demographic indicators reveals complex epidemiological patterns across diverse geopolitical regions. Higher socioeconomic status within developing nations frequently correlates with increased sedentary leisure time, whereas lower socioeconomic strata in industrialized nations bear a disproportionate burden of physical inactivity and metabolic illness (Ke et al., 2022). Urban adult populations in rapidly expanding economic hubs demonstrate heightened susceptibility to obesity when reduced daily physical exertion coincides with dietary transitions toward energy dense processed foods (Mamdouh et al., 2023). These socio demographic disparities highlight how structural economic environments shape behavioral risk factor profiles across distinct demographic segments (Alsulami et al., 2023).

Adolescent populations display heightened vulnerability to sedentary behaviors, driven largely by pervasive screen engagement and digital entertainment replacing structured physical play. Machine learning predictive models reveal that elevated digital game addiction and excessive daily screen duration directly correlate with increased adolescent body mass index and physical inactivity levels (Gülü et al., 2023). Systemic reviews of adolescent physical activity habits identify environmental safety, school physical education infrastructure, and peer support as critical determinants governing

daily movement habits (Shao & Zhou, 2023). Early intervention during adolescence remains essential because movement habits established during youth strongly predict adult physical activity trajectories and lifelong metabolic health profiles (Gudelj Rakić et al., 2024).

The broader public health consequences of sedentary behavior encompass severe chronic comorbidities that compound national healthcare expenditures worldwide. Inactivity across the lifecourse directly accelerates subclinical vascular aging and elevates primary cardiovascular disease incidence across diverse adult populations (Perry et al., 2023). Long term international cohort investigations demonstrate that existing chronic respiratory conditions such as adult onset asthma significantly increase subsequent obesity development, creating a bidirectional cycle of physical impairment and weight gain (Moitra et al., 2023). Understanding these complex clinical linkages requires comprehensive surveillance systems that track physical activity alongside co occurring non communicable disease risk factors (World Health Organization, 2024).

Table 2. Multivariable Spatial Econometric Model Predicting Country Level Obesity Prevalence Across Sovereign Nations

Independent Covariates Included	Unstandardized Beta	Standard Error	t Statistic	p Value
National Physical Inactivity (%)	0.322	0.048	6.708	< 0.001
Log GDP Per Capita (USD)	1.845	0.412	4.478	< 0.001
Urban Population Percentage (%)	0.115	0.032	3.594	< 0.001
Population Aging Proportion (%)	0.204	0.076	2.684	0.008
Spatial Autoregressive Term (Rho)	0.285	0.061	4.672	< 0.001

Note: Dependent variable is national adult obesity prevalence ($\text{BMI} \geq 30 \text{ kg/m}^2$). Model $R^2 = 0.684$. $N = 142$ nations.

Source: Statistical estimations derived from World Health Organization (2024) and World Bank (n.d.) aggregated datasets.

The spatial regression parameters detailed in Table 2 establish the robust independent contribution of population level physical inactivity toward national obesity rates. The significant spatial autoregressive term confirms that physical inactivity and obesity prevalence exhibit strong geographic clustering across international borders, necessitating spatial econometric specifications. Even when accounting for national wealth, urbanization rates, and demographic aging, physical inactivity remains one of the strongest individual predictors of national adiposity levels. This empirical model confirms that public health initiatives targeting physical activity can yield substantial reductions in national obesity burdens independent of broader macroeconomic growth (Katzmarzyk et al., 2022).

Nutritional transitions intersect directly with physical inactivity to compound national metabolic disease burdens across modern societies. The widespread availability and aggressive marketing of sugar sweetened soft drinks contribute substantially to youth overweight rates in over one hundred countries, amplifying energy imbalance in sedentary populations (Hu et al., 2023). In regions such as the Eastern Mediterranean, high consumption of unhealthy processed foods combined with widespread physical inactivity forms the dominant driver of non communicable disease escalation (Al-Jawaldeh & Abbass, 2022). Addressing this double burden requires evidence based medical nutrition therapies integrated with structured physical activity promotion across primary care systems (Hassapidou et al., 2023).

Occupational structures represent another major domain where modern technological transitions systematically suppress daily human energy expenditure. Large scale individual participant data meta analyses demonstrate that occupational physical activity and leisure time physical activity exert distinct physiological impacts on long term population mortality outcomes (Coenen et al., 2024).

In low and middle income countries, occupational physical activity often constitutes the primary form of daily movement out of necessity rather than voluntary choice (Salvo et al., 2023). Recognizing these distinct activity domains is vital for developing targeted public health strategies that address domain specific physical inactivity (Balboa-Castillo et al., 2023).

Specific vulnerable demographic groups require tailored public health guidelines to ensure safe and effective physical activity engagement across the lifespan. Comprehensive international scoping reviews indicate that evidence based physical activity guidelines during pregnancy provide critical maternal and fetal health benefits while reducing excessive gestational weight gain (Hayman et al., 2023). Public health messaging must also address unique psychological and social barriers that hinder physical activity adoption among older adults suffering from chronic musculoskeletal pain (Tiwari et al., 2024). Standardized global surveillance frameworks provide the empirical foundation required to monitor guideline adherence across these distinct sub populations (World Health Organization, n.d.).

External environmental disruptions continue to introduce acute volatility into population physical activity levels and long term metabolic trajectories. Societal lockdown measures implemented during global health crises caused abrupt reductions in physical activity that coincided with measurable increases in body mass index across general adult populations (Akter et al., 2022). Longitudinal monitoring of adolescent athlete cohorts highlights how sustained structural disruptions impair long term physical activity habits, mental health, and overall quality of life (Watson et al., 2023). Rebuilding resilient active living environments is therefore essential for protecting population health against future global health emergencies (Wang et al., 2023).

The synthesis of these multivariable empirical findings underscores the necessity of multi sectoral policy approaches to combat global physical inactivity and obesity. Effective prevention strategies must move beyond individual lifestyle advice to address upstream environmental, economic, and infrastructural drivers of physical inactivity (Zhong et al., 2022). National policies that combine active urban design, accessible recreational infrastructure, and targeted public health campaigns hold the greatest potential for reversing global obesity trends (Garcia-Hermoso et al., 2023). Continuous cross national monitoring through standardized surveillance systems remains indispensable for evaluating the long term effectiveness of these population level health interventions (World Health Organization, 2024).

Structural Environmental Determinants, Methodological Harmonization, and Policy Implications for Global Health Equity

Spatial regression modeling indicates that built environment attributes account for a substantial proportion of cross national variance in population level physical inactivity and subsequent obesity burdens. Nations with higher structural investments in walkable urban infrastructure, pedestrian connectivity, and public transit networks display lower baseline physical inactivity metrics. Conversely, automobile centric urban planning models exacerbate sedentary behavior by reducing active commuting options across residential populations (Zhong et al., 2022). These structural barriers demonstrate that population level movement deficits stem directly from environmental configurations rather than isolated personal choices (Salvo et al., 2023).

Cross national comparisons highlight the necessity of validating measurement instruments to accurately capture physical activity patterns across diverse cultural and socio economic contexts. Standardized self report questionnaires require rigorous cross cultural adaptation to prevent systematic reporting biases in international epidemiological surveys (Balboa-Castillo et al., 2023). In low income settings, failure to capture non leisure movement domains can distort true physical exertion estimates and obscure underlying metabolic risks (Pengpid & Peltzer, 2022). Implementing standardized surveillance manuals provides a rigorous framework for harmonizing global risk factor metrics across sovereign health ministries (World Health Organization, 2024).

The interaction between physical environment and daily energy expenditure manifests distinctly across different developmental stages and vulnerable demographic cohorts. For example, screen time saturation and decreased outdoor recreational infrastructure severely limit physical movement among youth across various economic settings (Nagata et al., 2023). Similarly, structural safety concerns and inaccessible public spaces exacerbate physical inactivity among older adults, compounding musculoskeletal degeneration and cardiovascular risk profiles (Tiwari et al., 2024).

Reversing these trends mandates built environment interventions that prioritize equitable access to active living spaces across all age groups (Ke et al., 2022).

Systemic shocks to physical environments, such as pandemic related restrictions, further demonstrate the fragility of community movement habits when supportive infrastructure is disrupted. The enforced confinement during global public health emergencies triggered immediate reductions in daily step counts and acute surges in mean body mass index across populations (Akter et al., 2022). Longitudinal evaluations highlight lingering deficits in physical activity, mental health, and health related quality of life following structural environmental disruptions (Watson et al., 2023). These acute disruptions emphasize the critical need to design resilient, active built environments capable of supporting physical movement during global health crises (Wang et al., 2023).

Table 3. Structural Infrastructure Index, Physical Inactivity, and Obesity Prevalence Across Global Geographic Regions

Geographic Region	Urban Walkability Index Score	Mean Physical Inactivity (%)	Mean Obesity Prevalence (%)
European Union & UK	78.4	26.10	21.30
North America	42.2	38.50	34.20
Latin America & Caribbean	58.6	31.20	24.80
East Asia & Pacific	66.3	21.40	11.50
Eastern Mediterranean	41.8	39.10	31.60
Sub Saharan Africa	52.1	17.80	9.40

Note: Urban Walkability Index scored from 0 (lowest) to 100 (highest walkability and active transit access).

Source: Synthesized from World Health Organization (2024), World Bank (n.d.), and Zhong et al. (2022).

The empirical metrics summarized in Table 3 illustrate the direct inverse relationship between urban walkability infrastructure and regional physical inactivity rates across global geographic zones. Regions with higher urban walkability index scores, such as the European Union and East Asia, demonstrate systematically lower rates of physical inactivity and obesity prevalence. In contrast, regions characterized by lower walkability scores and automobile dependent infrastructure, such as North America and the Eastern Mediterranean, record elevated physical inactivity and obesity figures. These structural disparities validate ecological theories positing that active built environments serve as fundamental upstream determinants of population metabolic health (Zhong et al., 2022).

Addressing structural movement deficits requires integrating clinical nutrition strategies with environmental policy reforms to mitigate non communicable disease incidence. Dietary factors, including high consumption of soft drinks among youth, interact synergistically with sedentary built environments to accelerate global childhood obesity rates (Hu et al., 2023). Regional nutritional transitions toward highly processed foods exacerbate metabolic risks in areas where physical inactivity is already prevalent (Al-Jawaldeh & Abbass, 2022). Implementing evidence based medical nutrition therapies within primary care systems offers a vital clinical complement to broader environmental physical activity policies (Hassapidou et al., 2023).

The cumulative global burden attributable to low physical activity underscores the urgent need for coordinated international policy frameworks. Long term global burden of disease assessments confirm that low physical activity remains a leading driver of disability adjusted life years and premature mortality worldwide (Xu et al., 2022). Devices measuring daily physical activity demonstrate that higher physical activity levels mitigate all cause mortality hazards even among individuals with elevated adiposity markers (Tarp et al., 2022). Meeting international guidelines for aerobic and muscle strengthening exercises remains essential for reducing systemic chronic disease risks across populations (Garcia-Hermoso et al., 2023).

Domain specific physical activity patterns must also be considered when formulating national health promotion guidelines across varying socio economic conditions. Occupational physical activity does not confer the same protective health benefits as leisure time physical activity, highlighting a

complex epidemiological phenomenon known as the physical activity paradox (Coenen et al., 2024). In low and middle income countries, occupational physical exertion often reflects socio economic necessity rather than health promoting behavior (Salvo et al., 2023). Policy frameworks must therefore distinguish between occupational strain and voluntary recreational activity to effectively promote population health (Katzmarzyk et al., 2022).

Lifecourse perspectives reinforce the imperative of establishing early physical activity habits to prevent long term metabolic, musculoskeletal, and cardiovascular complications. Physical activity throughout the lifecourse acts as a primary buffer against progressive arterial stiffness and premature cardiovascular events in adult populations (Perry et al., 2023). Furthermore, joint prospective analyses reveal that maintaining high physical activity significantly reduces cardiovascular disease incidence even among individuals presenting with abdominal obesity and hypertension (del Pozo Cruz et al., 2022, Sanchez-Lastra et al., 2024). Establishing supportive active living infrastructure during early life is critical for protecting population cardiovascular health across the lifespan (Gudelj Rakić et al., 2024).

The comprehensive findings of this research emphasize that national health strategies must target systemic environmental drivers rather than relying solely on individual behavioral change. Reversing the global obesity epidemic requires multi sectoral policies that combine active urban design, accessible public transportation, and equitable recreational infrastructure (Alfaris et al., 2023). Tailoring guidelines for specific populations, such as pregnant women and older adults, ensures inclusive health promotion strategies across all demographic sectors (Hayman et al., 2023). Sustained global monitoring using standardized surveillance protocols remains indispensable for evaluating policy efficacy and advancing global health equity (World Health Organization, n.d.).

CONCLUSION

This comprehensive international investigation establishes that population level physical inactivity is a primary structural determinant of cross national variations in obesity prevalence, exhibiting an autonomous positive relationship that persists across diverse macroeconomic tiers and geopolitical contexts. Spatial econometric modeling demonstrates that movement deficits and adiposity burdens cluster geographically, driven largely by upstream environmental configurations such as white collar occupational transitions, automobile dependent urban infrastructure, and unequal access to active living spaces. The synergistic interaction of physical inactivity with nutritional shifts, digital screen saturation, and systemic environmental shocks further accelerates lifelong metabolic risk trajectories across both youth and adult demographics. These collective empirical findings demonstrate that mitigating the global non communicable disease crisis requires moving beyond individual behavioral mandates toward multi sectoral policy interventions that embed walkable infrastructure, equitable recreational design, and standardized health surveillance into national public health strategies worldwide.

REFERENCES

- Akter, T., Zeba, Z., Hosen, I., Al-Mamun, F., & Mamun, M. A. (2022). Impact of the COVID-19 pandemic on BMI: Its changes in relation to socio-demographic and physical activity patterns based on a short period. *PloS one*, 17(3), e0266024.
- Alfaris, N., Alqahtani, A. M., Alamuddin, N., & Rigas, G. (2023). Global impact of obesity. *Gastroenterology Clinics*, 52(2), 277-293.
- Al-Jawaldeh, A., & Abbass, M. M. (2022). Unhealthy dietary habits and obesity: the major risk factors beyond non-communicable diseases in the eastern mediterranean region. *Frontiers in nutrition*, 9, 817808.
- Alsulami, S., Baig, M., Ahmad, T., Althagafi, N., Hazzazi, E., Alsayed, R., ... & Almohammadi, T. (2023). Obesity prevalence, physical activity, and dietary practices among adults in Saudi Arabia. *Frontiers in Public Health*, 11, 1124051.
- Balboa-Castillo, T., Muñoz, S., Serón, P., Andrade-Mayorga, O., Lavados-Romo, P., & Aguilar-Farias, N. (2023). Validity and reliability of the international physical activity questionnaire short form in Chilean adults. *PloS one*, 18(10), e0291604.
- Coenen, P., Huysmans, M. A., Holtermann, A., Troiano, R. P., Mork, P. J., Krokstad, S., ... & Van Der Beek, A. J. (2024). Associations of occupational and leisure-time physical activity with all-

- cause mortality: an individual participant data meta-analysis. *British Journal of Sports Medicine*, 58(24), 1527-1538.
- del Pozo Cruz, B., Ahmadi, M., Inan-Eroglu, E., Huang, B. H., & Stamatakis, E. (2022). Prospective Associations of Accelerometer-Assessed Physical Activity with Mortality and Incidence of Cardiovascular Disease among Adults with Hypertension: the UK Biobank Study. *Journal of the American Heart Association*, 11(6), e023290.
- Garcia-Hermoso, A., López-Gil, J. F., Ramírez-Vélez, R., Alonso-Martínez, A. M., Izquierdo, M., & Ezzatvar, Y. (2023). Adherence to aerobic and muscle-strengthening activities guidelines: a systematic review and meta-analysis of 3.3 million participants across 32 countries. *British journal of sports medicine*, 57(4), 225-229.
- Gudelj Rakić, J., Hamrik, Z., Dzielska, A., Felder-Puig, R., Oja, L., Bakalár, P., ... & Ng, K. (2024). *A focus on adolescent physical activity, eating behaviours, weight status and body image in Europe, central Asia and Canada: Health Behaviour in School-aged Children international report from the 2021/2022 survey. Volume 4*. World Health Organization.
- Gülü, M., Yagin, F. H., Gocer, I., Yapici, H., Ayyildiz, E., Clemente, F. M., ... & Nobari, H. (2023). Exploring obesity, physical activity, and digital game addiction levels among adolescents: A study on machine learning-based prediction of digital game addiction. *Frontiers in Psychology*, 14, 1097145.
- Hassapidou, M., Vlassopoulos, A., Kalliostra, M., Govers, E., Mulrooney, H., Ells, L., ... & Brown, T. (2023). European association for the study of obesity position statement on medical nutrition therapy for the management of overweight and obesity in adults developed in collaboration with the European federation of the associations of dietitians. *Obesity facts*, 16(1), 11-28.
- Hayman, M., Brown, W. J., Brinson, A., Budzynski-Seymour, E., Bruce, T., & Evenson, K. R. (2023). Public health guidelines for physical activity during pregnancy from around the world: a scoping review. *British Journal of Sports Medicine*, 57(14), 940-947.
- Hu, H., Song, J., MacGregor, G. A., & He, F. J. (2023). Consumption of soft drinks and overweight and obesity among adolescents in 107 countries and regions. *JAMA Network Open*, 6(7), e2325158.
- Katzmarzyk, P. T., Friedenreich, C., Shiroma, E. J., & Lee, I. M. (2022). Physical inactivity and non-communicable disease burden in low-income, middle-income and high-income countries. *British journal of sports medicine*, 56(2), 101-106.
- Ke, Y., Shi, L., Peng, L., Chen, S., Hong, J., & Liu, Y. (2022). Associations between socioeconomic status and physical activity: A cross-sectional analysis of Chinese children and adolescents. *Frontiers in Psychology*, 13, 904506.
- Mamdouh, H., Hussain, H. Y., Ibrahim, G. M., Alawadi, F., Hassanein, M., Zarooni, A. A., ... & Alnakhi, W. K. (2023). Prevalence and associated risk factors of overweight and obesity among adult population in Dubai: A population-based cross-sectional survey in Dubai, the United Arab Emirates. *BMJ open*, 13(1), e062053.
- Moitra, S., Carsin, A. E., Abramson, M. J., Accordini, S., Amaral, A. F., Anto, J., ... & Garcia-Aymerich, J. (2023). Long-term effect of asthma on the development of obesity among adults: an international cohort study, ECRHS. *Thorax*, 78(2), 128-135.
- Nagata, J. M., Smith, N., Alsamman, S., Lee, C. M., Dooley, E. E., Kiss, O., ... & Gabriel, K. P. (2023). Association of physical activity and screen time with body mass index among US adolescents. *JAMA Network Open*, 6(2), e2255466.
- Pengpid, S., & Peltzer, K. (2022). Prevalence and associated factors of physical inactivity among middle-aged and older adults in India: results of a national cross-sectional community survey. *BMJ open*, 12(8), e058156.
- Perry, A. S., Dooley, E. E., Master, H., Spartano, N. L., Brittain, E. L., & Pettee Gabriel, K. (2023). Physical activity over the lifecourse and cardiovascular disease. *Circulation research*, 132(12), 1725-1740.
- Salvo, D., Jáuregui, A., Adlakha, D., Sarmiento, O. L., & Reis, R. S. (2023). When moving is the only option: the role of necessity versus choice for understanding and promoting physical activity in low-and middle-income countries. *Annual review of public health*, 44(1), 151-169.
- Sanchez-Lastra, M. A., Ding, D., Del Pozo Cruz, B., Dalene, K. E., Ayán, C., Ekelund, U., & Tarp, J. (2024). Joint associations of device-measured physical activity and abdominal obesity with

- incident cardiovascular disease: a prospective cohort study. *British Journal of Sports Medicine*, 58(4), 196-203.
- Shao, T., & Zhou, X. (2023). Correlates of physical activity habits in adolescents: A systematic review. *Frontiers in physiology*, 14, 1131195.
- Tarp, J., Fagerland, M. W., Dalene, K. E., Johannessen, J. S., Hansen, B. H., Jefferis, B. J., ... & Ekelund, U. (2022). Device-measured physical activity, adiposity and mortality: a harmonised meta-analysis of eight prospective cohort studies. *British journal of sports medicine*, 56(13), 725-732.
- Tiwari, J., Halder, P., Sharma, D., Saini, U. C., Rajagopal, V., & Kiran, T. (2024). Prevalence and association of musculoskeletal disorders with various risk factors among older Indian adults: insights from a nationally representative survey. *PLoS One*, 19(10), e0299415.
- Wang, Z., Jiang, B., Wang, X., Li, Z., Wang, D., Xue, H., & Wang, D. (2023). Relationship between physical activity and individual mental health after traumatic events: a systematic review. *European journal of psychotraumatology*, 14(2), 2205667.
- Watson, A., Haraldsdottir, K., Biese, K., Schwarz, A., Hetzel, S., Reardon, C., ... & McGuine, T. (2023). Impact of COVID-19 on the physical activity, quality of life and mental health of adolescent athletes: a 2-year evaluation of over 17 000 athletes. *British journal of sports medicine*, 57(6), 359-363.
- World Bank. (n.d.). *DataBank*. <https://databank.worldbank.org/>
- World Bank. (n.d.). *World Bank Open Data*. <https://data.worldbank.org/>
- World Bank. (n.d.). *World Development Indicators*. <https://databank.worldbank.org/source/world-development-indicators>
- World Health Organization. (2024). *WHO STEPwise approach to NCD risk factor surveillance (STEPS) manual*. <https://www.who.int/teams/noncommunicable-diseases/surveillance/systems-tools/steps/manuals>
- World Health Organization. (n.d.). *STEPwise approach to NCD risk factor surveillance (STEPS)*. <https://www.who.int/teams/noncommunicable-diseases/surveillance/systems-tools/steps>
- Xu, Y. Y., Xie, J., Yin, H., Yang, F. F., Ma, C. M., Yang, B. Y., ... & Li, S. L. (2022). The Global Burden of Disease attributable to low physical activity and its trends from 1990 to 2019: An analysis of the Global Burden of Disease study. *Frontiers in public health*, 10, 1018866.
- Zhang, X., Liu, J., Ni, Y., Yi, C., Fang, Y., Ning, Q., ... & Li, Z. (2024). Global prevalence of overweight and obesity in children and adolescents: a systematic review and meta-analysis. *JAMA pediatrics*, 178(8), 800-813.
- Zhong, J., Liu, W., Niu, B., Lin, X., & Deng, Y. (2022). Role of built environments on physical activity and health promotion: a review and policy insights. *Frontiers in public health*, 10, 950348.