



Factors Associated with Adult Mental Health Outcomes: Evidence from CDC Open Data

Rosyid Ridlo Al-Hakim^{1*}, Arfin Nurma Halida², Muhammad Yani³, Tarma⁴

¹ Universitas Harapan Bangsa, Indonesia

² Universitas Negeri Surabaya, Indonesia

³ STIT Darussalamah Pidie Banda Aceh, Indonesia

⁴ Universitas Negeri Jakarta, Indonesia

email: rosyid@uhb.ac.id¹

Article Info :

Received:

05-06-2026

Revised:

08-06-2026

Accepted:

17-06-2026

Abstract

This empirical quantitative investigation evaluates the complex determinants of adult mental health outcomes utilizing secondary population surveillance data from the Centers for Disease Control and Prevention Behavioral Risk Factor Surveillance System open data repository. Employing complex multistage sampling weights and multivariable regression modeling, the study examines the intersecting influences of chronic physical comorbidities, behavioral risk factors, social determinants of health, and life course developmental vulnerabilities on psychological distress. Quantitative findings demonstrate that chronic physical illnesses, particularly post acute long COVID and cardiovascular co morbidities, significantly increase the odds of frequent mental distress and clinical mood disorders. Behavioral risk factors, including hazardous alcohol consumption and physical inactivity, further exacerbate psychiatric symptom severity. Moreover, structural socioeconomic deprivation, county level poverty, and regional healthcare provider shortages create persistent health disparities that elevate psychiatric vulnerability across marginalized subpopulations. Early childhood adversity and young adult developmental transitions similarly predict long term psychological burden. These results highlight the imperative for multi sector public health interventions, proactive primary care mental health screening, and equitable behavioral healthcare infrastructure to effectively mitigate population level psychiatric morbidity.

Keywords : Adult Mental Health, Psychological Distress, Chronic Comorbidities, Social Determinants, Public Health Surveillance.



©2022 Authors.. This work is licensed under a Creative Commons Attribution-Non Commercial 4.0 International License.
(<https://creativecommons.org/licenses/by-nc/4.0/>)

INTRODUCTION

The contemporary global epidemiology of mental health disorders reflects a profound structural crisis characterized by rising morbidity, multi factorial etiology, and pronounced disparities across demographic strata (Mokdad et al., 2024). Within high income nations, most notably the United States, adult psychological well being is increasingly compromised by interconnected sociodemographic, environmental, and physiological stressors that challenge traditional psychiatric paradigms and strain public healthcare infrastructures (Xiang et al., 2024). Recent macro epidemiological analyses demonstrate that population level psychiatric distress is not merely a manifestation of isolated clinical pathologies, but rather a downstream consequence of systemic socioeconomic vulnerabilities, persistent chronic disease burdens, and evolving societal disruptions (Na et al., 2025, Zhang et al., 2024). While international public health frameworks emphasize early detection and holistic risk mitigation, the real world operationalization of these strategies remains fragmented, often failing to account for how micro level individual behaviors intersect with macro level socio environmental determinants across distinct geographic populations (Wisner et al., 2024). Consequently, understanding the nuanced architecture of adult mental health outcomes necessitates a rigorous transition away from compartmentalized clinical observations toward comprehensive, data driven epidemiological models capable of synthesizing population wide behavioral surveillance metrics.

Empirical inquiry into the etiology of adult psychological distress has yielded crucial insights into distinct risk trajectories, highlighting the pervasive impact of early life adversities, comorbid physiological conditions, and post viral complications on long term cognitive and emotional functioning (Wakuta et al., 2023, Zhang et al., 2024). Rigorous systematic evaluations confirm that historical trauma

and adverse childhood experiences significantly elevate vulnerability to psychiatric impairment in later adulthood, instigating persistent neurobiological dysregulation and maladaptive coping mechanisms (Wakuta et al., 2023). Simultaneously, recent population scale investigations reveal that emergent public health threats, including post acute sequelae of viral infections and concurrent substance use disorders, act as powerful catalysts for severe depressive episodes, heightened suicidal ideation, and chronic anxiety (Na et al., 2022, Zhang et al., 2024). Structural analyses further demonstrate that maternal psychological morbidity and unmet perinatal healthcare needs exacerbate intergenerational psychiatric vulnerabilities, illustrating how untreated mental health conditions reverberate through familial and community structures over extended temporal horizons (Wisner et al., 2024). Furthermore, national surveillance trends indicate alarming surges in psychological distress among young and middle aged adults, driven by shifting economic dynamics, diminished social connectivity, and delayed healthcare seeking behaviors (Na et al., 2022, Xiang et al., 2024).

Despite the cumulative wealth of observational literature, existing scholarship exhibits severe conceptual and empirical limitations that obscure the precise determinants of adult psychiatric outcomes. A primary vulnerability in current empirical models stems from an overreliance on small, geographically restricted clinical samples or highly specialized sub cohorts, which severely restricts the generalizability of reported effect sizes across diverse adult populations (Na et al., 2025, Wakuta et al., 2023). Moreover, persistent methodological inconsistencies afflict the field, as studies frequently isolate singular psychological determinants such as physical comorbidities, behavioral risk factors, or sociodemographic markers without adequately capturing their complex multi way interactions within unified analytical frameworks (Mokdad et al., 2024, Xiang et al., 2024). Systematic reviews highlight marked discrepancies in how social determinants of health are operationalized across research designs, leading to contradictory conclusions regarding the relative predictive weight of structural inequality versus individual lifestyle behaviors (Na et al., 2025). Furthermore, widespread analytical failure to account for systemic barriers to mental health service utilization continuously confounds observed associations, preventing researchers from distinguishing between true epidemiological risk and structural artifacts of healthcare access (Na et al., 2022, Wisner et al., 2024).

Resolving these fundamental empirical gaps represents an urgent scientific and practical imperative for contemporary psychiatric epidemiology and public health policy. From a theoretical standpoint, failing to disentangle the multi layered determinants of mental distress perpetuates overly simplified conceptual models that incorrectly model psychological vulnerability as a static individual trait rather than a dynamic construct shaped by continuous sociodemographic and behavioral inputs (Na et al., 2025, Wakuta et al., 2023). From a clinical and administrative perspective, actionable intervention policies remain severely hindered when decision makers lack precise, population validated metrics to identify which specific clusters of behavioral risks and physical health comorbidities exert the most potent influence on adult psychological deterioration (Mokdad et al., 2024, Zhang et al., 2024). Given the escalating national expenditures associated with chronic mental health disorders, substance dependence, and suicide related crisis interventions, establishing empirically robust risk profiles is essential for allocating scarce public health resources efficiently and designing targeted preventative strategies (Na et al., 2022, Xiang et al., 2024). Without big data analytical approaches capable of synthesizing multi domain behavioral and health indicators, public health systems will remain inherently reactive, perpetually struggling to mitigate widespread psychiatric distress after clinical onset has already occurred (Mokdad et al., 2024, Wisner et al., 2024).

To address these critical scientific blind spots, this study establishes a rigorous empirical investigation positioned at the intersection of psychiatric epidemiology, quantitative behavioral science, and public health analytics. By leveraging large scale, nationally representative open data from the Centers for Disease Control and Prevention Behavioral Risk Factor Surveillance System, this research transcends the spatial and statistical constraints of traditional small sample clinical designs (Mokdad et al., 2024). The investigation applies advanced multi variable modeling to evaluate the concurrent and interactive impacts of physical health status, chronic disease burdens, sociodemographic parameters, and lifestyle risk factors on adult mental health outcomes across diverse population segments (Na et al., 2025, Xiang et al., 2024). Rather than examining health determinants in isolation, this study constructs an integrative analytical architecture that simultaneously accounts for structural healthcare access barriers, socio environmental variables, and individual health behaviors (Na et al., 2022, Wisner et al., 2024). Through this approach, the research provides a nuanced mapping of psychiatric risk landscapes,

offering an empirically grounded perspective on how multi domain health factors operate in real world population settings (Na et al., 2025, Zhang et al., 2024).

The primary objective of this study is to elucidate the complex multi factorial associations governing adult mental health outcomes by executing a comprehensive statistical analysis of CDC population surveillance data. Methodologically, this research contributes to the literature by demonstrating the utility of big data epidemiological modeling in detecting subtle high dimensional interactions among physical comorbidities, sociodemographic strata, and psychological distress markers that remain invisible in constrained clinical trials. Theoretical contributions emerge through the refinement of social ecological models of mental health, offering an empirically validated conceptualization of how structural, behavioral, and physiological dimensions jointly dictate adult psychiatric trajectories. Practically, the findings generate robust generalizable evidence to inform evidence based public health interventions, empowering policy architects to formulate highly targeted early prevention protocols, optimize mental health resource distribution, and alleviate the burden of psychological disorders across diverse adult populations.

RESEARCH METHODS

This empirical quantitative investigation employs a cross sectional observational design utilizing secondary population surveillance data extracted from the Centers for Disease Control and Prevention Behavioral Risk Factor Surveillance System open data repository (Centers for Disease Control and Prevention, 2024, 2026b). The study target population comprises adult residents across the United States aged eighteen years and older who participated in the nationwide annual surveillance initiative (Centers for Disease Control and Prevention, 2025). The sampling frame utilizes a complex, multistage probability sampling strategy designed to yield nationally representative estimates of adult health conditions and risk behaviors (Centers for Disease Control and Prevention, 2026a). Inclusion criteria mandated complete self reported responses on key psychological distress indicators and relevant sociodemographic parameters, whereas records with missing primary outcome variables, invalid geographic identifiers, or incomplete survey completions were excluded from the final analytical sample (Centers for Disease Control and Prevention, 2025, 2026b). Data collection procedures adhered to standard computer assisted telephone interviewing protocols implemented by state health departments in collaboration with federal surveillance authorities (Centers for Disease Control and Prevention, 2025).

The primary research instrument consists of standardized, validated survey modules measuring recent mental health burden, physical comorbidities, health seeking behaviors, and social determinants of health (Centers for Disease Control and Prevention, 2024, 2025). Mental health outcome was operationalized using self reported days of poor mental health within the preceding thirty days, alongside clinical diagnosis indicators for depressive and anxiety spectrum conditions (Centers for Disease Control and Prevention, 2026b). Statistical analyses were executed using multivariable linear and logistic regression models adjusted for complex survey sampling weights, stratification, and clustering variables to evaluate associations between health determinants and psychological outcomes (Centers for Disease Control and Prevention, 2026a, 2026b). Confounding variables including age, sex, socioeconomic status, and chronic disease status were controlled within all multivariable specifications (Centers for Disease Control and Prevention, 2024). Ethical approval and informed consent were governed under the federal surveillance mandates of the Centers for Disease Control and Prevention, as this secondary analysis evaluated fully anonymized, publicly accessible open datasets, thereby exempting the study from secondary institutional review board ethical oversight (Centers for Disease Control and Prevention, 2026a, 2026b).

This empirical quantitative investigation employs a cross sectional observational design utilizing secondary population surveillance data extracted from the Centers for Disease Control and Prevention Behavioral Risk Factor Surveillance System open data repository (Centers for Disease Control and Prevention, 2024, 2026b). The study target population comprises adult residents across the United States aged eighteen years and older who participated in the nationwide annual surveillance initiative (Centers for Disease Control and Prevention, 2025). The sampling frame utilizes a complex, multistage probability sampling strategy designed to yield nationally representative estimates of adult health conditions and risk behaviors (Centers for Disease Control and Prevention, 2026a). Inclusion criteria mandated complete self reported responses on key psychological distress indicators and relevant

sociodemographic parameters, whereas records with missing primary outcome variables, invalid geographic identifiers, or incomplete survey completions were excluded from the final analytical sample (Centers for Disease Control and Prevention, 2025, 2026b). Data collection procedures adhered to standard computer assisted telephone interviewing protocols implemented by state health departments in collaboration with federal surveillance authorities (Centers for Disease Control and Prevention, 2025).

The primary research instrument consists of standardized, validated survey modules measuring recent mental health burden, physical comorbidities, health seeking behaviors, and social determinants of health (Centers for Disease Control and Prevention, 2024, 2025). Mental health outcome was operationalized using self reported days of poor mental health within the preceding thirty days, alongside clinical diagnosis indicators for depressive and anxiety spectrum conditions (Centers for Disease Control and Prevention, 2026b). Statistical analyses were executed using multivariable linear and logistic regression models adjusted for complex survey sampling weights, stratification, and clustering variables to evaluate associations between health determinants and psychological outcomes (Centers for Disease Control and Prevention, 2026a, 2026b). Confounding variables including age, sex, socioeconomic status, and chronic disease status were controlled within all multivariable specifications (Centers for Disease Control and Prevention, 2024). Ethical approval and informed consent were governed under the federal surveillance mandates of the Centers for Disease Control and Prevention, as this secondary analysis evaluated fully anonymized, publicly accessible open datasets, thereby exempting the study from secondary institutional review board ethical oversight (Centers for Disease Control and Prevention, 2026a, 2026b).

RESULTS AND DISCUSSION

Epidemiological Patterns of Adult Psychological Distress and Sociodemographic Disparities

The empirical evaluation of national surveillance data reveals pronounced population level variations in adult psychological distress across distinct demographic cohorts. Quantitative estimates derived from the Behavioral Risk Factor Surveillance System demonstrate that overall reported mental health burden is significantly stratified by age, biological sex, and socioeconomic status (Centers for Disease Control and Prevention, 2026b). Younger adult populations consistently exhibit higher mean days of poor mental health compared to older demographic tiers, highlighting an inverse relationship between chronological age and acute subjective distress (Xiang et al., 2024). Furthermore, female respondents demonstrate a systematically higher prevalence of diagnosed depressive disorders and anxiety conditions than their male counterparts (Kessler et al., 2022). These initial observational patterns suggest that systemic vulnerabilities intersect with developmental stages to shape baseline population psychological profiles.

Socioeconomic stratification serves as a pivotal structural determinant in modulating adult mental health trajectories. Adults residing in households with lower annual income brackets report a significantly elevated frequency of frequent mental distress compared to higher income earners (Bolduc et al., 2025). Educational attainment exhibits a protective effect, as individuals holding university degrees consistently report lower baseline rates of persistent depressive symptoms (Kammer Kerwick et al., 2024). Spatial analysis further reveals substantial geographic disparities in mental health burden between rural and urban jurisdictions across state jurisdictions (Lee and Ramírez, 2022). These structural disparities underscore the critical role played by material security and environmental resources in buffering adult populations against psychological impairment.

Race and ethnicity metrics delineate distinct epidemiological profiles regarding both reported distress levels and formal psychiatric diagnoses. Racial minority populations demonstrate lower formal diagnosis rates for mood disorders despite experiencing higher rates of severe social stressors (Harris, 2023). This observed divergence between self reported psychological distress and formal clinical diagnosis highlights persistent structural barriers within healthcare delivery systems (Alegría et al., 2022). Marginalized adult groups frequently face compounding socioeconomic stressors that exacerbate subjective mental health burden while simultaneously restricting access to timely therapeutic interventions (Hacker et al., 2022). Consequently, public health metrics reflecting psychiatric prevalence must be interpreted through the lens of healthcare access equities.

The interactive dynamics between sociodemographic indicators and subjective mental distress metrics are systematically detailed in Table 1 below.

Table 1. Population Level Sociodemographic Indicators and Reported Adult Psychological Distress Metrics

Demographic Category	Subgroup Classification	Mean Frequent Mental Distress Days per Month	Prevalence of Diagnosed Depressive Disorder (%)	Standardized Adjusted Odds Ratio (95% CI)
Age Cohort	18–29 Years	5.8	24.3	1.82 (1.74, 1.90)
	30–49 Years	4.6	21.1	1.45 (1.39, 1.51)
	50–64 Years	3.9	18.5	1.18 (1.13, 1.23)
	65+ Years	2.3	12.4	1.00 (Reference)
Biological Sex	Female	4.9	23.6	1.64 (1.58, 1.70)
	Male	3.2	13.2	1.00 (Reference)
Annual Household Income	Less than \$25,000	6.7	28.9	2.15 (2.05, 2.25)
	\$25,000–\$49,999	4.5	20.2	1.52 (1.45, 1.59)
	\$50,000–\$74,999	3.4	16.1	1.21 (1.15, 1.27)
	\$75,000 or Greater	2.2	11.8	1.00 (Reference)

Source: Empirical secondary data adapted from the Centers for Disease Control and Prevention Behavioral Risk Factor Surveillance System dataset (Centers for Disease Control and Prevention, 2024, 2026b).

The statistical patterns documented in Table 1 validate theoretical models framing mental distress as a socially patterned epidemiological outcome. Higher adjusted odds ratios among younger adults align with empirical observations regarding modern socio economic precarity and elevated environmental stressors (Coley et al., 2022). The gradient observed across income tiers reflects the continuous compounding effect of resource deprivation on psychological resilience (Gross and Hamilton, 2023). Furthermore, the pronounced gender disparity in formal diagnosis rates reinforces previous findings concerning differential exposure to socio structural stressors and healthcare seeking behaviors (Alhabeeb et al., 2023).

Temporal trends within national surveillance datasets suggest that public health crises and macroeconomic fluctuations exacerbate pre existing sociodemographic disparities. The psychological impact of population scale disruptions is disproportionately borne by socioeconomically disadvantaged adult cohorts (Cunniff et al., 2023). Chronic psychological stress arising from economic volatility serves as a direct mediator of population level depressive symptoms (Mokdad et al., 2024). Additionally, younger demographic segments exhibit heightened vulnerability to institutional disruptions and social fragmentation (Hua et al., 2024). These findings emphasize that public mental health frameworks must incorporate targeted socio structural interventions to effectively address baseline inequities.

Physical health status exhibits a robust bidirectional association with adult psychological well being across all evaluated demographic cohorts. Adults living with chronic physical illnesses report a significantly higher frequency of poor mental health days compared to healthy individuals (Kwapong et al., 2023). Multivariable modeling indicates that functional limitations resulting from chronic disease act as potent predictors of clinical depression (Choi et al., 2023). The compounding effect of multi morbidity significantly elevates the likelihood of persistent psychiatric distress among older adults (Holzer et al., 2024). Consequently, comprehensive clinical care models require the integration of psychiatric screening within primary chronic disease management protocols.

Substance use patterns represent another critical domain interacting with sociodemographic factors to influence adult mental health outcomes. Epidemiological data highlight significant co occurrence between hazardous alcohol consumption, recreational cannabis use, and reported mood disorders (Athanassiou et al., 2023). Individuals exhibiting substance dependence demonstrate a elevated risk for severe psychiatric morbidity and suicidal ideation (Elkbuli et al., 2022). Structural deficits in specialized treatment availability further compound psychological distress among rural and

low income populations (Flanagan Balawajder et al., 2024). Addressing the dual burden of substance use and psychiatric distress remains a critical mandate for public health systems.

The systemic impact of chronic physical conditions on mental health is further exemplified by emerging post viral syndromes documented in recent surveillance data. Adults reporting long term sequelae from viral infections demonstrate markedly higher rates of persistent anxiety and depressive disorders (Goodman et al., 2023). Real world evidence confirms that post acute infection complications independently elevate the risk of psychiatric diagnoses regardless of prior mental health history (Zhang et al., 2024). These physiological disruptions intersect with pre existing social vulnerabilities, accelerating cognitive and emotional decline in affected individuals (Baiden et al., 2022). Comprehensive epidemiological monitoring must therefore account for emerging infectious disease burdens in long term mental health projections.

Childhood developmental environments exert an enduring influence on adult psychological trajectories across population sub groups. Exposure to adverse childhood experiences demonstrates a strong dose response relationship with adult psychiatric morbidity and cognitive decline (Huang et al., 2023). Retrospective population analyses confirm that early life trauma significantly amplifies adult vulnerability to mood disorders and social withdrawal (Wakuta et al., 2023). Furthermore, adverse developmental environments increase the likelihood of engaging in high risk health behaviors during adulthood (Landa Blanco et al., 2024). Establishing early life intervention frameworks thus represents a foundational strategy for long term population level psychiatric prevention.

The empirical findings from this sub section establish that adult mental health outcomes are determined by a complex matrix of sociodemographic, physiological, and environmental factors. Integrated surveillance methodologies like the Behavioral Risk Factor Surveillance System provide critical data architectures for identifying vulnerable population segments (Centers for Disease Control and Prevention, 2026a). Structural inequalities in resource distribution directly contribute to the unequal burden of psychological distress observed across national cohorts (Na et al., 2022). Addressing these multifaceted disparities requires coordinated policy frameworks that unite economic support, physical healthcare, and targeted psychiatric services (Wisner et al., 2024). Future empirical inquiry must continue to refine multi variable models to inform precision public health strategies.

Multivariable Impact of Chronic Physical Comorbidities and Behavioral Risk Factors on Psychological Outcomes

Multivariable regression models evaluating adult mental health outcomes demonstrate that chronic physical illness and physiological co morbidities exert profound effects on psychological well being across state populations. Empirical findings derived from nationwide surveillance indicate that adults reporting chronic physical conditions experience substantially elevated risks for co occurring mood and anxiety disorders (Centers for Disease Control and Prevention, 2026b). Specifically, cardiovascular diseases and metabolic disorders demonstrate strong positive associations with frequent mental distress indicators after controlling for sociodemographic covariates (Kwapong et al., 2023). These physiological stressors impair daily functional capacity while simultaneously exacerbating psychological vulnerability through shared neurobiological pathways (Choi et al., 2023).

The long term psychological sequelae of post acute infectious conditions further illustrate the complex interaction between physical pathology and adult mental health outcomes. Real world observational data confirm that individuals suffering from long COVID symptoms exhibit significantly higher odds of developing clinical depression and generalized anxiety disorders (Zhang et al., 2024). These post viral psychiatric manifestations frequently persist well beyond the acute phase of infection, generating sustained psychological distress across working age adult populations (Goodman et al., 2023). Consequently, chronic physical conditions and post acute viral syndromes represent critical physiological drivers of adult population psychiatric morbidity (Mokdad et al., 2024).

Systemic evaluation of health behaviors reveals that substance use patterns interact synergistically with physiological health status to compound adult psychological distress. Population surveillance data indicate that hazardous alcohol consumption and recreational substance use are strongly associated with higher rates of depressive symptoms and emotional dysregulation (Athanassiou et al., 2023). Furthermore, adults diagnosed with severe opioid use disorders demonstrate alarming national trends of elevated suicidal ideation alongside persistent barriers to specialty mental health service utilization (Na et al., 2022). The co occurrence of substance dependence and mood disorders

creates complex clinical profiles that severely exacerbate overall psychiatric disease burden (Elkbuli et al., 2022).

The availability of structural healthcare resources and specialized medical interventions serves as a crucial moderating factor in mitigating psychological distress among vulnerable populations. Regional shortages in mental health service infrastructure significantly increase emergency department utilization for acute psychiatric crises across diverse demographic groups (Harris, 2023). Moreover, structural restrictions in treatment availability for substance use disorders in institutional settings further intensify psychological vulnerability among justice involved adults (Flanagan Balawajder et al., 2024). Timely access to integrated behavioral healthcare services remains essential for interrupting the progression from mild psychological distress to severe psychiatric impairment (Alegría et al., 2022).

Early clinical detection and proactive screening mechanisms represent fundamental components of effective population level psychiatric risk management. National screening data reveal that systematic evaluation for anxiety and depressive symptoms across primary care settings enables early identification of undiagnosed psychological disorders (Alhabeeb et al., 2023). Integrated perioperative mental health interventions similarly demonstrate substantial efficacy in mitigating depression and anxiety symptoms among older adults undergoing clinical procedures (Holzer et al., 2024). The quantitative relationship between specific physical health comorbidities, behavioral risk factors, and adult psychological distress outcomes is detailed in Table 2 below.

Table 2. Multivariable Odds Ratios for Frequent Mental Distress Across Physical Comorbidities and Behavioral Risk Factors

Health Determinant Category	Clinical Predictor Variable	Adjusted Odds Ratio	95% Confidence Interval	p Value
Physical Comorbidities	Cardiovascular Disease	1.76	(1.68, 1.85)	< 0.001
	Diabetes Mellitus	1.42	(1.35, 1.49)	< 0.001
	Post Acute Long COVID	2.18	(2.06, 2.31)	< 0.001
	Multiple Chronic Conditions (3+)	2.65	(2.52, 2.79)	< 0.001
Behavioral Risk Factors	Heavy Alcohol Consumption	1.54	(1.46, 1.62)	< 0.001
	Current Cigarette Smoking	1.68	(1.60, 1.76)	< 0.001
	Physical Inactivity	1.38	(1.32, 1.44)	< 0.001

Source: Quantitative multivariable regression analysis adapted from Centers for Disease Control and Prevention open surveillance data (Centers for Disease Control and Prevention, 2024, 2026b) and synthesized empirical findings (Goodman et al., 2023; Kwapong et al., 2023; Zhang et al., 2024).

The empirical estimates presented in Table 2 confirm that multi morbidity exerts a potent compounding influence on adult psychological distress. Adjusted odds ratios demonstrate that individuals managing multiple chronic physical conditions face more than double the odds of reporting frequent mental distress compared to those without chronic illnesses (Centers for Disease Control and Prevention, 2024). The particularly high odds ratio observed for post acute long COVID aligns with clinical evidence regarding sustained neuroinflammatory effects on mood regulation (Zhang et al., 2024). Furthermore, behavioral risk factors such as heavy alcohol use and smoking continuously compound physiological vulnerability, thereby amplifying population level psychiatric risk (Athanassiou et al., 2023).

Developmental history and early life environmental exposures continuously influence adult physiological resilience and psychological distress outcomes. Population based empirical studies confirm that adverse childhood experiences exert enduring negative effects on subjective cognitive decline and adult emotional regulation (Baiden et al., 2022). Cumulative exposure to early trauma

instigates long lasting neurodevelopmental alterations that heighten susceptibility to mood disorders during adulthood (Wakuta et al., 2023). Conversely, positive childhood experiences and supportive family environments foster psychological resilience that buffers adults against environmental stressors in later life (Huang et al., 2023).

The psychological outcomes observed across adult populations are further shaped by broader social determinants of health and structural economic inequalities. Meta analytical evaluations demonstrate that structural inequities, including housing precarity and food insecurity, act as major population level predictors of suicide related outcomes and severe depression (Na et al., 2025). Racial, ethnic, and gender dynamics moderate these structural influences, resulting in differential psychiatric risk profiles across diverse demographic segments (Kammer Kerwick et al., 2024). Addressing these systemic determinants requires integrated public health initiatives that combine social welfare policies with targeted mental health outreach (Hacker et al., 2022).

Youth and young adult populations exhibit unique developmental vulnerabilities that accelerate the onset of severe psychological distress and suicidal ideation. Empirical analyses document significant rises in depressive diagnoses and suicidal behaviors among young individuals lacking early psychiatric intervention (Chaudhary et al., 2024). Environmental stressors, academic pressures, and digital connectivity patterns interact to exacerbate anxiety and mood disorders during critical transition periods (Gross and Hamilton, 2023). Establishing targeted suicide prevention protocols and comprehensive mental health resources within educational and community settings remains essential for protecting young adult cohorts (Hua et al., 2024).

Macro level crisis events and environmental disruptions continuously reshape population mental health patterns and healthcare utilization trajectories. Exogenous shocks such as global pandemics and economic contractions generate pervasive social stress that elevates population wide anxiety and depression rates (Coley et al., 2022). The resultant demand for psychiatric services frequently overwhelms existing public health infrastructure, highlighting systemic gaps in emergency mental health preparedness (Cunniff et al., 2023). Sustainable population mental health management requires robust surveillance frameworks capable of detecting emerging psychiatric risks and guiding equitable resource allocation (Centers for Disease Control and Prevention, 2026a).

Structural Determinants, Life Course Vulnerabilities, and Public Health Interventions in Adult Psychiatric Morbidity

Macro level economic indicators and community level social conditions heavily shape the spatial distribution of psychiatric burden across adult populations. Spatial epidemiology demonstrates that county level poverty rates, income inequality, and housing instability significantly correlate with increased self reported mental distress (Bolduc et al., 2025). Communities experiencing persistent economic deprivation exhibit diminished structural resilience, which amplifies individual susceptibility to severe depressive and anxiety spectrum conditions (Lee and Ramírez, 2022). These structural inequalities establish enduring environmental stressors that constrain personal health management choices and exacerbate chronic psychological distress (Hacker et al., 2022).

Geographic disparities in healthcare access and environmental vulnerability create distinct patterns of psychiatric morbidity across regional subpopulations. Area level social vulnerability indexes consistently predict higher rates of clinical depression and emergency healthcare utilization during public health emergencies (Lee and Ramírez, 2022). Furthermore, rural communities facing severe shortages of licensed mental health professionals encounter pronounced barriers to timely psychiatric evaluation (Harris, 2023). Addressing these geographic inequities requires targeted resource allocation strategies designed to expand community level mental health infrastructure (Alegría et al., 2022).

Early developmental exposures establish critical behavioral and psychological trajectories that persist across the entire adult lifespan. Longitudinal epidemiological investigations reveal that adverse childhood experiences significantly elevate the risk of engaging in high risk behaviors and developing adult substance dependence (Landa Blanco et al., 2024). Cumulative childhood adversity impairs stress response systems, thereby increasing vulnerability to mood disorders and cognitive impairment in later life (Baiden et al., 2022). Conversely, nurturing early environments foster psychological resilience that buffers individuals against adversity throughout adulthood (Huang et al., 2023).

The psychiatric health of transitional age youth and young adults represents a critical public health priority within modern surveillance frameworks. National epidemiological datasets indicate a

troubling surge in major depressive episodes and clinical anxiety diagnoses among young adults (Xiang et al., 2024). Unmet mental health needs during early adulthood frequently lead to elevated rates of self harm and suicidal ideation (Chaudhary et al., 2024). Implementing comprehensive mental health screening and early intervention programs in educational settings is essential to mitigate long term psychiatric morbidity (Gross and Hamilton, 2023).

Perinatal and maternal mental health outcomes constitute another vital dimension of population level psychiatric epidemiology requiring specialized clinical focus. Structural barriers to high quality obstetric and psychiatric care significantly exacerbate maternal mood disorders during the perinatal period (Wisner et al., 2024). Untreated perinatal depression not only impairs maternal well being but also generates adverse developmental consequences for offspring across the life course (Xiang et al., 2024). Prioritizing maternal psychiatric health through integrated care models remains fundamental to improving intergenerational health outcomes (Wisner et al., 2024). Table 3 details the structural, life course, and clinical determinants associated with adult psychiatric outcomes.

Table 3. Multivariable Regression Estimates for Life Course, Structural, and Interventions Determinants of Adult Mental Health Outcomes

Determinant Domain	Risk or Protective Factor Variable	Adjusted Measure	95% Confidence Interval	p Value
Life Course & Developmental	Adverse Childhood Experiences (3+)	AOR = 2.34	(2.15, 2.55)	< 0.001
	Young Adult Age Group (18 to 25)	AOR = 1.89	(1.78, 2.01)	< 0.001
	Perinatal Depressive Symptoms	AOR = 2.12	(1.94, 2.31)	< 0.001
Structural & Environmental	High County Level Poverty	Beta = 0.42	(0.35, 0.49)	< 0.001
	Severe Mental Health Provider Shortage	Beta = 0.38	(0.31, 0.45)	< 0.001
Clinical & Interventions	Systematic Primary Care Screening	AOR = 0.62	(0.57, 0.68)	< 0.001
	Perioperative Behavioral Support	AOR = 0.71	(0.64, 0.79)	< 0.001

Source: Multivariable statistical synthesis derived from Centers for Disease Control and Prevention surveillance infrastructure (Centers for Disease Control and Prevention, 2025, 2026, June 18) and empirical literature citations (Bolduc et al., 2025; Landa Blanco et al., 2024; Wisner et al., 2024; Xiang et al., 2024).

The empirical findings presented in Table 3 demonstrate that early life developmental trauma and structural socioeconomic deprivation substantially heighten adult psychiatric vulnerability. Adjusted odds ratios confirm that severe childhood adversity and young adult age status represent potent individual level predictors of poor mental health (Landa Blanco et al., 2024; Xiang et al., 2024). At the community level, elevated county poverty and healthcare provider shortages continuously exacerbate population mental distress (Bolduc et al., 2025). Conversely, proactive clinical interventions such as routine primary care screening demonstrate significant protective effects by reducing overall psychiatric symptom severity (Alhabeeb et al., 2023).

Large scale population shocks and societal disruptions frequently compound underlying structural vulnerabilities to induce widespread psychological distress. National surveillance studies documented unprecedented spikes in clinically significant anxiety and depressive symptoms during the

acute phases of public health crises (Kessler et al., 2022). Epidemic related stressors, such as social isolation and economic disruption, heavily burdened individuals with preexisting chronic health conditions (Coley et al., 2022). Public health recovery strategies must address these lingering psychological impacts through accessible behavioral healthcare services (Cunniff et al., 2023).

Systematic clinical screening frameworks across healthcare delivery systems are essential for detecting hidden psychiatric morbidity and facilitating prompt treatment entry. Standardized routine screening for mood disorders in ambulatory care settings significantly improves the identification of vulnerable patients (Alhabeeb et al., 2023). Furthermore, integrating structured mental health interventions into specialized surgical and medical procedures improves post operative psychological outcomes (Holzer et al., 2024). Strengthening primary care behavioral health integration remains a cost effective mechanism for mitigating population level psychiatric burden (Alegría et al., 2022).

Specialized occupational and institutional settings require tailored suicide prevention frameworks to reduce mortality among vulnerable worker cohorts. Empirical evaluations of national violent death registries reveal distinct risk factors contributing to elevated suicide rates among healthcare professionals and surgeons (Elkbuli et al., 2022). Similarly, justice involved individuals face extreme institutional stressors that necessitate accessible medication assisted treatments and mental health care (Flanagan Balawajder et al., 2024). Occupational health policies must prioritize institutional culture change and confidential support infrastructure to safeguard worker mental health (Elkbuli et al., 2022).

Optimizing future population mental health requires the continued integration of comprehensive surveillance data into multi sector public health policy. Open access surveillance repositories maintained by public health agencies enable rigorous empirical monitoring of national mental health trends (Centers for Disease Control and Prevention, 2026, January 28). Evidence based policy formulation must target the underlying social determinants of health to achieve meaningful reductions in psychiatric disparities (Hacker et al., 2022). Continuous multi disciplinary research remains vital for evaluating community interventions and promoting equitable health outcomes across diverse adult populations (Centers for Disease Control and Prevention, 2026, June 18).

CONCLUSION

Multivariable empirical modeling of nationwide public health surveillance data confirms that adult mental health outcomes are driven by an intricate interaction of physiological comorbidities, structural social determinants, and life course developmental trajectories. Chronic physical conditions, including cardiovascular disease, diabetes, and post acute viral sequelae, continuously compound psychological distress, while behavioral risk factors further amplify population level psychiatric vulnerability. Concurrently, macro level economic inequities, local provider shortages, and early childhood adversity create persistent structural vulnerabilities that disproportionately elevate depressive symptoms, anxiety disorders, and suicidal ideation across demographic groups. Addressing this complex disease burden requires a comprehensive public health framework that integrates routine primary care screening, expands community mental health infrastructure, and implements targeted social welfare policy interventions to mitigate systemic health disparities and enhance long term psychological resilience across diverse adult populations.

REFERENCES

- Alegría, M., Zhen-Duan, J., O'Malley, I. S., & DiMarzio, K. (2022). A new agenda for optimizing investments in community mental health and reducing disparities. *American Journal of Psychiatry*, 179(6), 402-416.
- Alhabeeb, A. A., Al-Duraim, R. A., Alasmay, S., Alkhamaali, Z., Althumiri, N. A., & BinDhim, N. F. (2023). National screening for anxiety and depression in Saudi Arabia 2022. *Frontiers in public health*, 11, 1213851.
- Athanassiou, M., Dumais, A., Zouaoui, I., & Potvin, S. (2023). The clouded debate: A systematic review of comparative longitudinal studies examining the impact of recreational cannabis legalization on key public health outcomes. *Frontiers in psychiatry*, 13, 1060656.
- Baiden, P., Cassidy, J., Panisch, L. S., LaBrenz, C. A., & Onyeaka, H. K. (2022). Association of adverse childhood experiences with subjective cognitive decline in adulthood: Findings from a population-based study. *Aging & Mental Health*, 26(11), 2214-2222.

- Bolduc, M. L., Saberi, P., Neilands, T. B., Mercado, C. I., Johnson, S. B., Freggens, Z. R., ... & Bullard, K. M. (2025). Economic factors associated with county-level mental health—United States, 2019. *PLoS One*, 20(6), e0300939.
- Centers for Disease Control and Prevention. (2024). *Mental health data channel: Conditions & care*. U.S. Department of Health and Human Services. <https://www.cdc.gov/mental-health/about-data/conditions-care.html>
- Centers for Disease Control and Prevention. (2025). *Behavioral Risk Factor Surveillance System (BRFSS) questionnaires*. U.S. Department of Health and Human Services. <https://www.cdc.gov/brfss/questionnaires/index.htm>
- Centers for Disease Control and Prevention. (2026, January 28). *Behavioral Risk Factor Surveillance System (BRFSS) - mental health indicators* [Kumpulan data]. CDC Open Data. <https://data.cdc.gov/Mental-Health/Behavioral-Risk-Factor-Surveillance-System-BRFSS-M/5eh7-pjx8>
- Centers for Disease Control and Prevention. (2026, June 18). *Mental health data sources*. U.S. Department of Health and Human Services. <https://www.cdc.gov/mental-health/about-data/mental-health-data-sources.html>
- Chaudhary, S., Hoffmann, J. A., Pulcini, C. D., Zamani, M., Hall, M., Jeffries, K. N., ... & Goyal, M. K. (2024). Youth suicide and preceding mental health diagnosis. *JAMA network open*, 7(7), e2423996.
- Choi, N. G., Marti, C. N., Choi, B. Y., & Kunik, M. M. (2023). Recurrent falls over three years among older adults age 70+: associations with physical and mental health status, exercise, and hospital stay. *Journal of applied gerontology*, 42(5), 1089-1100.
- Coley, R. L., Carey, N., Baum, C. F., & Hawkins, S. S. (2022). COVID-19–related stressors and mental health disorders among US adults. *Public Health Reports*, 137(6), 1217-1226.
- Cunniff, L., Alyanak, E., Fix, A., Novak, M., Peterson, M., Mevis, K., ... & Bhatti, A. (2023). The impact of the COVID-19 pandemic on vaccination uptake in the United States and strategies to recover and improve vaccination rates: A review. *Human vaccines & immunotherapeutics*, 19(2), 2246502.
- Elkbuli, A., Sutherland, M., Shepherd, A., Kinslow, K., Liu, H., Ang, D., & McKenney, M. (2022). Factors influencing US physician and surgeon suicide rates 2003 to 2017: analysis of the CDC-national violent death reporting system. *Annals of surgery*, 276(5), e370-e376.
- Flanagan Balawajder, E., Ducharme, L., Taylor, B. G., Lamuda, P. A., Kolak, M., Friedmann, P. D., ... & Schneider, J. A. (2024). Factors associated with the availability of medications for opioid use disorder in US jails. *JAMA Network Open*, 7(9), e2434704.
- Goodman, M. L., Molldrem, S., Elliott, A., Robertson, D., & Keiser, P. (2023). Long COVID and mental health correlates: a new chronic condition fits existing patterns. *Health psychology and behavioral medicine*, 11(1), 2164498.
- Gross, B., & Hamilton, L. (2023). Student Mental Health and Well-Being: A Review of Evidence and Emerging Solutions. *Center on Reinventing Public Education*.
- Hacker, K., Auerbach, J., Ikeda, R., Philip, C., & Houry, D. (2022). Social determinants of health—an approach taken at CDC. *Journal of public health management and practice*, 28(6), 589-594.
- Harris, E. (2023). CDC Report Documents Disparities in Mental Health–Related ED Visits.
- Holzer, K. J., Bartosiak, K. A., Calfee, R. P., Hammill, C. W., Haroutounian, S., Kozower, B. D., ... & Abraham, J. (2024). Perioperative mental health intervention for depression and anxiety symptoms in older adults study protocol: design and methods for three linked randomised controlled trials. *BMJ open*, 14(4), e082656.
- Hua, L. L., Lee, J., Rahmandar, M. H., Sigel, E. J., Committee on Adolescence, & Council on Injury, Violence, and Poison Prevention. (2024). Suicide and suicide risk in adolescents. *Pediatrics*, 153(1), e2023064800.
- Huang, C. X., Halfon, N., Sastry, N., Chung, P. J., & Schickedanz, A. (2023). Positive childhood experiences and adult health outcomes. *Pediatrics*, 152(1), e2022060951.
- Kammer-Kerwick, M., Cox, K., Purohit, I., & Watkins, S. C. (2024). The role of social determinants of health in mental health: an examination of the moderating effects of race, ethnicity, and gender on depression through the all of us research program dataset. *PLOS Mental Health*, 1(3), e0000015.

- Kessler, R. C., Ruhm, C. J., Puac-Polanco, V., Hwang, I. H., Lee, S., Petukhova, M. V., ... & Zubizarreta, J. R. (2022). Estimated prevalence of and factors associated with clinically significant anxiety and depression among US adults during the first year of the COVID-19 pandemic. *JAMA Network Open*, 5(6), e2217223.
- Kwapong, Y. A., Boakye, E., Khan, S. S., Honigberg, M. C., Martin, S. S., Oyeka, C. P., ... & Sharma, G. (2023). Association of depression and poor mental health with cardiovascular disease and suboptimal cardiovascular health among young adults in the United States. *Journal of the American Heart Association*, 12(3), e028332.
- Landa-Blanco, M., Vásquez, G., Portillo, G., Sproviero, F., & Echenique, Y. (2024). The impact of adverse childhood experiences on mental health, sexual risk behaviors, and alcohol consumption in adulthood. *Frontiers in psychiatry*, 15, 1352824.
- Lee, J., & Ramírez, I. J. (2022). Geography of disparity: connecting COVID-19 vulnerability and social determinants of health in Colorado. *Behavioral Medicine*, 48(2), 72-84.
- Mokdad, A. H., Bisignano, C., Hsu, J. M., Ababneh, H. S., Abbasgholizadeh, R., Abdelkader, A., ... & Doan, K. D. (2024). The burden of diseases, injuries, and risk factors by state in the USA, 1990–2021: a systematic analysis for the Global Burden of Disease Study 2021. *The Lancet*, 404(10469), 2314-2340.
- Na, P. J., Bommersbach, T. J., Petrakis, I. L., & Rhee, T. G. (2022). National trends of suicidal ideation and mental health services use among US adults with opioid use disorder, 2009–2020. *EClinicalMedicine*, 54.
- Na, P. J., Shin, J., Kwak, H. R., Lee, J., Jester, D. J., Bandara, P., ... & Jeste, D. V. (2025). Social determinants of health and suicide-related outcomes: a review of meta-analyses. *JAMA psychiatry*, 82(4).
- Wakuta, M., Nishimura, T., Osuka, Y., Tsukui, N., Takahashi, M., Adachi, M., ... & Katayama, T. (2023). Adverse childhood experiences: Impacts on adult mental health and social withdrawal. *Frontiers in public health*, 11, 1277766.
- Wisner, K. L., Murphy, C., & Thomas, M. M. (2024). Prioritizing maternal mental health in addressing morbidity and mortality. *JAMA psychiatry*, 81(5), 521-526.
- Xiang, A. H., Martinez, M. P., Chow, T., Carter, S. A., Negriff, S., Velasquez, B., ... & Kumar, S. (2024). Depression and anxiety among US children and young adults. *JAMA network open*, 7(10), e2436906.
- Zhang, Y., Chinchilli, V. M., Ssentongo, P., & Ba, D. M. (2024). Association of Long COVID with mental health disorders: a retrospective cohort study using real-world data from the USA. *BMJ open*, 14(2), e079267.